

On the rotation of polarization by a gravitational lens

Valerio Faraoni

Department of Physics and Astronomy, University of Victoria

P.O. Box 3055, Victoria, B.C. V8W 3P6 (Canada)

Abstract

It is proved that the field of a gravitational lens induces no rotation in the polarization vector of electromagnetic radiation, in agreement with the previous literature, but with a different approach. The result is generalized to the case of less conventional gravitational lenses (static cosmic strings and gravitational waves).

To appear in *Astronomy and Astrophysics*

1 Introduction

The problem of propagation of polarized electromagnetic radiation through the gravitational field of a cosmic object is accompanied by a rotation of the plane of polarization of the radiation, has been considered recently (Dyer & Shaver 1992, hereafter DS). If such an effect would exist, it could be relevant for the analysis of photons coming from a gravitationally lensed source, or for the microwave background photons propagating in an inhomogeneous universe. It has been proposed (Kronberg et al. 1991) that polarization data from gravitationally lensed radio sources can be used to probe the mass distribution in the lens. In the paper by Kronberg et al. (1991), it is assumed that the polarization vector of the electromagnetic radiation is not rotated by the gravitational lens. This assumption has been discussed in a later paper by DS, who have proved it for most astrophysically interesting lenses, mainly using symmetry arguments in a relativistic context. Our purpose is to compare the result by DS with an alternative approach, adopting a post-Newtonian metric to describe spacetime, and a Newtonian potential to describe the gravitational field of the lens, which are more conventional tools in gravitational lens theory. Calculations are performed in Sec. 2, and the result by DS is confirmed. In Sec. 3, we consider the case of more exotic gravitational lenses which have been treated in the literature, namely cosmic strings and gravitational waves. It is shown that the negative result of Sec. 2 holds also for these nonconventional lenses. A symmetry argument analogous to those in DS' paper is used for the case of a static, straight, infinite cosmic string, while for gravitational waves the calculation is closely analogous to that for ordinary gravitational lenses.

2 Gravitational lenses and polarized radiation

Let us consider a radio source which is lensed by a gravitational lens described by the Newtonian potential ϕ ; we assume that

1. the spacetime is described by the post-Newtonian metric¹

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} , \quad (2.1)$$

where $\eta_{\mu\nu}$ are the components of the Minkowski metric in an asymptotically Cartesian coordinate system $\{x^\mu\} = \{t, x, y, z\}$, and

$$h_{00} = -2\phi , \quad (2.2)$$

¹The metric signature is +2. Greek indices run from 0 to 3, Latin indices run from 1 to 3. We use units in which $G = c = 1$, and perform computations to first order in ϕ and its derivatives.

$$h_{ij} = -2\phi \delta_{ij} , \quad (2.3)$$

are small perturbations;

2. geometric optics holds;
3. light rays suffer only small deflections;
4. the lens is bounded and stationary, i.e.

$$\phi(x, y, z) \rightarrow 0 , \quad \partial_i \phi \rightarrow 0 \quad \text{as} \quad r \equiv (x^2 + y^2 + z^2)^{1/2} \rightarrow +\infty ; \quad (2.4)$$

$$\frac{\partial \phi}{\partial t} \simeq 0 . \quad (2.5)$$

In the present calculations, we assume that the observer is an inertial observer of the Minkowski background, with four velocity components $u^\mu = \delta^{0\mu}$ in the $\{x^\alpha\}$ coordinate system. As customary in gravitational lens studies, cosmology can be fit into the model at a later stage, by using angular diameter distances in a Friedmann-Lemaitre-Robertson-Walker universe to measure the distances between observer, lens, and source. We will actually need to go beyond the geometric optics approximation, in order to describe polarization of the electromagnetic radiation.

An electromagnetic wave in a curved spacetime can be described by the Maxwell four-potential

$$A_\mu = \hat{A}_\mu(x^\alpha) e^{i\omega S(x^\alpha)} , \quad (2.6)$$

where $\hat{A}^\mu$ is a complex vector field, S is a real function (the eikonal), and ω is the frequency of the electromagnetic wave. A^μ and $S^\mu \equiv S^{\cdot\mu}$ satisfy (Stephani 1982)

$$S_\mu S^\mu = 0 , \quad (2.7)$$

$$A_\mu S^\mu = 0 , \quad (2.8)$$

$$S^\nu \nabla_\nu S^\mu = 0 . \quad (2.9)$$

The complex vector $\hat{A}^\mu$ can be decomposed into its complex magnitude a and the real polarization vector P^μ

$$\hat{A}^\mu = a P^\mu , \quad (2.10)$$

where a and P^μ obey the equations (Stephani 1982)

$$\frac{1}{a} \frac{da}{d\lambda} = -\theta , \quad (2.11)$$

$$\frac{dP^\mu}{d\lambda} = \frac{1}{2} \left(\frac{P^\nu \partial_\nu a}{a} + \nabla_\nu P^\nu \right) S^\mu . \quad (2.12)$$

Here, λ is an affine parameter along the null geodesics, and $\theta \equiv \nabla_\alpha S^\alpha/2$ is the expansion of the congruence of null geodesics around a fiducial ray, which obeys the well known Raychaudhuri's equation. We can write

$$a = a^{(0)} + \delta a , \quad (2.13)$$

$$P^\mu = P^{(0)\mu} + \delta P^\mu , \quad (2.14)$$

where δa and δP^μ are small perturbations to the flat space quantities $a^{(0)}$ and $P^{(0)\mu}$.

We choose, as a solution to Eqs. (2.11) and (2.12) in the unperturbed case (for which $\theta^{(0)} = 1/\lambda$),

$$a^{(0)} = \frac{A}{\lambda} , \quad (2.15)$$

$$P^{(0)\mu} = (0, 1, 0, 0) , \quad (2.16)$$

where A is a complex constant, the unperturbed photons are polarized along the x -axis, and the affine parameter $\lambda = t$ is measured from the source position. We assume now that the optic axis of the lens coincides with the z -axis, and consider a bundle of null rays such that the corresponding photons have unperturbed paths parallel to it, and are only slightly deflected by the gravitational lens. The four-vector tangent to a perturbed path is

$$S^\mu = S^{(0)\mu} + \delta S^\mu = (1 + \delta S^0, \delta S^1, \delta S^2, 1 + \delta S^3) , \quad (2.17)$$

where $S^{(0)\mu} \equiv (1, 0, 0, 1)$ is the tangent to the unperturbed path. The equation of null geodesics

$$\frac{dS^\mu}{d\lambda} + \Gamma_{\rho\sigma}^\mu S^\rho S^\sigma = 0 , \quad (2.18)$$

and

$$\Gamma_{\rho\sigma}^\mu = \frac{1}{2} \eta^{\mu\alpha} (h_{\alpha\rho,\sigma} + h_{\alpha\sigma,\rho} - h_{\rho\sigma,\alpha}) , \quad (2.19)$$

give the following expression for the deflection δS^μ

$$\delta S^\mu = - \int_S^O d\lambda \left(h_{\rho,\sigma}^\mu - \frac{1}{2} h_{\rho\sigma}{}^{,\mu} \right) S^{(0)\rho} S^{(0)\sigma} + O(2) , \quad (2.20)$$

where the integral in the right hand side is computed along the perturbed photon path from the source to the observer. Equation (2.20) gives, to first order,

$$\delta S^0 = 0 , \quad (2.21)$$

$$\delta S^i = -2 \int_S^O dz \partial_i \phi + O(2) , \quad (2.22)$$

$$\delta S^3 = 0 \quad (2.23)$$

($i = 1, 2$), where the integral in the right hand side of Eq. (2.22) is now computed along the *unperturbed* photon path from the source to the observer (performing the integration along the actual path only adds a second order contribution). The gravitational lens induces no frequency shifts and no deflections in the z -direction, to first order, as can be seen from Eqs. (2.21) and (2.23), and is well known.

The perturbations δa and δP^μ satisfy

$$\frac{1}{a^{(0)}} \frac{d(\delta a)}{d\lambda} + \frac{1}{\lambda} \frac{\delta a}{a^{(0)}} + \delta\theta = 0 , \quad (2.24)$$

$$\frac{d(\delta P^\mu)}{d\lambda} = \frac{1}{2} \left[\frac{P^{(0)\nu} \partial_\nu(\delta a)}{a^{(0)}} + \nabla_\nu P^\nu \right] S^\mu , \quad (2.25)$$

with $\delta\theta \equiv \theta - \theta^{(0)}$ given by the Raychaudhuri's equation. For our purposes, it is not necessary to solve the coupled system of Eqs. (2.24), (2.25), but it is sufficient to note that, using Eqs. (2.5), (2.15), (2.23), and

$$\sqrt{|g|} = 1 - 2\phi + O(2) , \quad (2.26)$$

Eq. (2.25) reduces to

$$\frac{d(\delta P^\alpha)}{d\lambda} = \frac{1}{2} \left[\frac{\partial_x(\delta a)}{a^{(0)}} + \partial_\mu(\delta P^\mu) - 2 \partial_x \phi \right] (\delta^{0\alpha} + \delta^{3\alpha}) . \quad (2.27)$$

We have, from Eq. (2.27),

$$\frac{d(\delta P^1)}{d\lambda} = \frac{d(\delta P^2)}{d\lambda} = 0 \quad (2.28)$$

and, since $\delta P^\mu \rightarrow 0$ as $r \rightarrow +\infty$ (at least approaching the source of radiation), we get

$$\delta P^1 = \delta P^2 = 0 . \quad (2.29)$$

Since P^μ is a purely spatial vector, $\delta P^0 = 0$. Moreover, Eq. (2.27) gives

$$\frac{d(\delta P^0)}{d\lambda} = \frac{d(\delta P^3)}{d\lambda} , \quad (2.30)$$

and we conclude that $\delta P^3 = \delta P^0 = 0$ as well. Thus, to first order, the gravitational lens induces no changes in the polarization vector of electromagnetic radiation:

$$\delta P^\mu = 0 , \quad (2.31)$$

in agreement with the result by DS.

3 Discussion and conclusion

The action of a gravitational lens on the plane of polarization of electromagnetic radiation from a distant source has been found to be negligible to first order, according to the result by DS. We describe the gravitational lens with a Newtonian potential, as customary in the conventional gravitational lens theory (Bourassa et al. 1973; Bourassa & Kantowski 1975; Schneider 1985; Blandford & Narayan 1986; Blandford & Kochanek 1988). This approach is alternative to that of DS, but it is obviously limited to situations in which the assumptions needed to apply the standard scalar or vector formalism (based on the use of a post-Newtonian metric) hold. In the case of rotating or, more generally, nonstationary gravitational lenses, these formalisms cannot be employed, or their use requires a particular care (Faraoni 1991). However, it seems that most objects of astrophysical interest whose action as gravitational lenses is likely to be observed, behave as static, and can therefore be described by the vector or scalar formalism. This class of objects include conventional lenses like galaxies, galaxy clusters, microlenses like stars, Jupiters, binary systems, and even compact objects like neutron stars or black holes, provided that lensing takes place in a region where the field is sufficiently weak (a situation much more more likely than lensing in a strong field region). The velocities of galaxies in clusters, of stars in galaxies, and the rotational velocities of double galaxies, or binary systems of stars, planets, or collapsed objects (provided that the system is not an extremely close binary), are not high enough to make the lens nonstationary (Faraoni 1991). The case of lensing by an extremely rotating lens should be considered as highly unlikely in the real world, as pointed out in DS. A Schwarzschild lens, though static, cannot be described by the scalar or vector formalism of gravitational lensing theory, except for the weak field regime. In the strong field region, the symmetry arguments of DS are particularly convenient.

In principle, one can also conceive of more exotic lenses. The gravitational field due to density perturbations which are collapsing to form structures in the early universe is by far too weak to affect appreciably the polarization of the microwave background photons propagating close to them. Another possibility is represented by lensing cosmic strings (Vilenkin 1984; Hogan & Narayan 1984; Paczynski 1986; Birkinshaw 1989). The spacetime around a straight, static, infinite string can be described by the metric

$$ds^2 = -dt^2 + dr^2 + (1 - 8\mu) r^2 d\varphi^2 + dz^2, \quad (3.1)$$

where μ is the linear mass density of the string. This metric can be brought to the Minkowskian form by introducing the coordinate $\varphi' \equiv (1 - 4\mu) \varphi$, which varies in the range $[0, (1 - 4\mu) 2\pi]$. Though spacetime is locally flat, the existence of a deficit angle

$\delta = 4\mu$ causes the deflection of a light ray propagating in the xy plane, and the formation of a double image of a distant source. By using symmetry arguments analogous to those of DS, one concludes that the polarization vector of the electromagnetic radiation is not affected by the string, since the latter does not introduce any preferred verse of rotation of the polarization vector (one can reach the same conclusion by using Eqs. (2.11) and (2.12) in the metric given by Eq. (3.1)). A possible exception is given by a cosmic string moving at relativistic velocity (Birkinshaw 1989); in this case the symmetry argument does not hold. However, no observational evidence has been given for such objects, and we will not investigate this situation here.

Another possibility considered in the literature is lensing by gravitational waves, both considered as lens components superposed to the caustic structure of an ordinary gravitational lens (McBreen & Metcalfe 1988; Allen 1989, 1990; Kovner 1990), or as full lenses (Wheeler 1960; Zipoy 1966; Zipoy & Bertotti 1968; Kaufmann 1970; Bertotti 1971; Dautcourt 1974; Bertotti and Catenacci 1975; Burke 1975; Linder 1986, 1988; Braginsky et al. 1990; Faraoni 1991, 1992). As far as these situations are concerned, we remind the reader that the shear induced by a gravitational wave creates an anisotropy in the plane orthogonal to direction of propagation of the wave. This results in a polarization of the order of the metric perturbations describing the gravitational wave; for astrophysically generated gravitational waves, this effect is too small to be detectable with present techniques. In the case of long wavelength gravitational waves of cosmological origin (which cannot be treated in the context of the thin lens approximation), the effect on the polarization of microwave background photons has been taken into account, and has been used to set upper limits on the cosmological density of such waves (Matzner 1988). However, this effect is like the photon scattering in an anisotropic medium, and is quite different from the rotation of the plane of polarization. In fact, assuming that the spacetime metric is given by

$$g_{\mu\nu} = \eta_{\mu\nu} + \gamma_{\mu\nu} , \quad (3.2)$$

where $\gamma_{\mu\nu}$ are small perturbations describing the gravitational waves, one finds, with calculations closely parallel to those in Sec. 2,

$$\frac{d(\delta P^\alpha)}{d\lambda} = \frac{1}{2} \left[\frac{\partial_x(\delta a)}{a^{(0)}} + \partial_\mu(\delta P^\mu) + \frac{1}{2} \partial_x \gamma \right] (\delta^{0\alpha} + \delta^{3\alpha}) , \quad (3.3)$$

where

$$\gamma \equiv \gamma^\mu{}_\mu = -\gamma_{00} + \gamma_{11} + \gamma_{22} + \gamma_{33} + O(2) . \quad (3.4)$$

Equation (3.3) is analogous to Eq. (2.27). The same arguments used above lead to the result that $\delta P^\mu = 0$ in this case as well.

As a conclusion, any common kind of gravitational lens does not induce rotation of polarization, at least at the level detectable by present techniques.

Acknowledgments

Doctor P. Schneider is acknowledged for helpful comments. The author would like to thank also the warm hospitality at the University of Victoria, in particular Professor F. I. Cooperstock. This work was supported by the “Fondazione Angelo della Riccia”, which is gratefully acknowledged.

References

- Allen, B. 1989, Phys. Rev. Lett. 63, 2017
 Allen, B. 1990, Gen. Relativ. Grav. 22, 1447
 Bertotti, B. 1971, in Sachs, R.K. (ed.) General Relativity and Cosmology. Academic Press, New York, p.347
 Bertotti, B., Catenacci, R. 1975, Gen. Relativ. Grav. 6, 329
 Birkinshaw, M. 1989, in Moran, J.M., Hewitt, J.N., Lo K.Y. (eds.) Gravitational Lenses, Proceedings Cambridge, Massachusetts, USA, 1988. Springer, Berlin, p. 59-64
 Blandford, R.D., Kochanek, C.S. 1988, in Bachall, J.N., Piran, T., Weinberg, S. (eds.) Proceedings of the 13th Jerusalem Winter School on Dark Matter in the Universe. World Scientific, Singapore, p. 133
 Blandford, R.D., Narayan, R. 1986, ApJ 310, 568
 Bourassa, R.R., Kantowski, R. 1975, ApJ 195, 13
 Bourassa, R.R., Kantowski, R., Norton, T.D. 1973, ApJ 185, 747
 Braginsky, V.B., Kardashev, N.S., Polnarev, A.G., Novikov, I.D. 1990, Nuovo Cimento 105B, 1141
 Burke, W.L. 1975, ApJ 196, 329
 Dautcourt, G. 1974, in Longair, M.S. (ed.) IAU Symp. 63, Confrontation of Cosmological Theories with Observation. Reidel, Dordrecht, p. 299
 Dyer C.C. Shaver, E.G. 1992, ApJ 390, L5 (DS)
 Faraoni, V. 1991, preprint SISSA 161/91/A, to appear in ApJ
 Faraoni, V. 1992, in Kayser, R., Schramm, T., Refsdal, S. (eds.) Gravitational Lenses, Proceedings, Hamburg, Germany 1991. Springer, Berlin (in press)

- Hogan, C., Narayan, R. 1984, MNRAS 211, 575
- Kaufmann, W.J. 1970, Nat 227, 157
- Kovner, I. 1990, ApJ 351, 114
- Kronberg, P.P., Dyer, C.C., Burbidge, E.M., Junkkarinen, V.T. 1991, ApJ 367, L1
- Linder, E.V. 1986, Phys. Rev. D 34, 1759
- Linder, E.V. 1988, ApJ 328, 77
- Matzner, R.A. 1988, Univ. of Texas preprint
- McBreen, B., Metcalfe, L. 1988, Nat 332, 234
- Paczynski, B. 1986, Nat 319, 567
- Schneider, P. 1985, A&A 143, 413
- Stephani, H. 1982, General Relativity: An Introduction to the Theory of the Gravitational Field. Cambridge Univ. Press, p.75-77
- Vilenkin, A. 1984, ApJ 282, L51
- Wheeler, J.A. 1960, in Radicati, L.A. (ed.) Rendiconti della Scuola Internazionale di Fisica “Enrico Fermi”, 11th Course of the Varenna Summer School, 1959, “Interazioni Deboli”. Zanichelli, Bologna, p.67
- Winterberg, F. 1968, Nuovo Cimento 53B, 195
- Zipoy, D.M. 1966, Phys. Rev. 142, 825
- Zipoy, D.M., Bertotti, B. 1968, Nuovo Cimento 56B, 195